

# The mystery of accelerating turbidity currents: The curious case of the Congo Canyon

A.J. Cotel<sup>1</sup> and R.E. Breidenthal<sup>2</sup>

<sup>1</sup>*Dept. Of Civil and Environmental Engineering, University of Michigan, Ann Arbor, MI 48109*

<sup>2</sup>*Department of Aeronautics and Astronautics, University of Washington Seattle, WA 98195*

*(Dated: September 10, 2020)*

Motivated by the remarkably large propagation distances observed in turbidity currents near the mouth of the Congo River in Africa, a new model is proposed for their dynamics. It assumes that the erosion of solid particles from the bed underneath the current increases the density of the current such that the vortex rotational rate increases over the case of no erosion. If the rate of increase of vortex rotation is sufficient, the entrainment rate of fluid above the current is inhibited. As a consequence, the turbidity current propagates much farther than would be expected without the dynamic effect of acceleration.

## I. INTRODUCTION

Originally discovered after the sequential failure of transatlantic cables, turbidity currents play a central role in the transport of sediment in the continental shelf and out into intro-continental basins, abyssal plains, and deep lakes. Turbidity currents are a type of two-phase, density currents. In other contexts, lahars, powder snow avalanches, pyroclastic, and lava flows also obey similar physics [1].

These currents can be highly destructive. For example, Mt. Rainier in Washington state is regarded by the US Geological Survey as the most dangerous volcano in the US, primarily because of the threat of fast-moving lahars to descend from the mountain into densely populated valleys near Seattle and Tacoma. Such flows need not be triggered by any volcanic eruption. A simple slope failure can suffice. Avalanches kill on average about 28 people every year in the U.S. [2]

The strongest known turbidity currents are in the Congo Canyon off the west coast of equatorial Africa [3], transporting sediment a great distance. Conventional values of turbulent entrainment rates suggest that these currents should more promptly slow down and stop due to entrainment of ambient fluid [4]. It appears that some unknown physics is inhibiting the turbulent entrainment.

## II. BACKGROUND

After some triggering event such as a slope failure, a mass of sediment descends a submarine slope. The mixture is more dense than the surrounding pure fluid, so it forms a density current flowing down the slope. Baroclinic torques generate vorticity, and the flow is certainly turbulent at typical Reynolds numbers of roughly  $10^6$ . The turbulent eddies entrain pure upper fluid, tending to dilute the mixture and reduce the density contrast. If the bed underneath the turbidity current is erodible, the current may incorporate additional sediment, tending to increase the density contrast. So there are two opposing effects that compete to alter the density difference.

The most important dimensionless parameter in buoyant flows is the Richardson number, the ratio of potential to kinetic energy. It is defined as follows:

$$Ri = g' \delta / W^2, \quad (1)$$

where

$$g' = g \Delta \rho / \rho \quad (2)$$


is the reduced acceleration of gravity,  $\Delta \rho$  is the density difference between the current and the ambient fluid,  $\rho$  is the density, and  $g$  is the acceleration of gravity. The characteristic thickness and speed of the current are  $\delta$  and  $W$  respectively. The entrainment velocity of the

---

<sup>1</sup> [acotel@umich.edu](mailto:acotel@umich.edu)

<sup>2</sup> [breident@aa.washington.edu](mailto:breident@aa.washington.edu)

ambient fluid from above the current is  $v_a$ , and the bed



erosion rate from below is  $v_b$ . Figure 1 is a sketch.

FIG. 1. Density current

Buoyant flows tend to adjust themselves to achieve a Richardson number of approximately one, corresponding to an equipartition of energy between potential and kinetic. The flow does this by adjusting the speed, entraining fluid, and/or by shedding fluid [5].

Entrainment plays the most important role in the dynamics of turbulent flows in general and of turbidity currents in particular. According to the entrainment hypothesis of Morton *et al.* [6], the entrainment velocity is always proportional to the characteristic velocity of the flow, equivalent to saying that the entrainment velocity depends only on the largest eddy size. The hypothesis makes perfect sense when the largest eddy is the one doing the entraining.

However, the hypothesis fails when entrainment occurs by an eddy size different from the largest. For example, in a stratified or compressible flow, the governing dimensionless parameter of Richardson or Mach number defines the size of the entraining eddy to be less than the largest. The entrainment hypothesis is no longer valid.

Just as the entrainment hypothesis fails when a smaller eddy does the entraining, it also fails under strong vortex acceleration. Here acceleration refers to the rotational rate of turbulent eddies. First observed in the turbulent jet [7], [8], and [9], the theory was extended to all canonical turbulent flows [10]. Counter-intuitively, the

normalized entrainment rate decreases in all accelerating flows, except Rayleigh-Taylor.

In particular, acceleration effects can play a large role in buoyant flows. The first known example is the heated jet, a laboratory simulation of buoyancy-addition in cumulus clouds due to latent heat from phase change [11]. They observed the spreading angle of an upward-moving vertical jet decrease as heat was added to it.

Note that the entrainment rate is reduced even when the rotation rate of the large eddy is still decreasing in time. What matters is the rate of decrease with respect to that of the unaccelerated flow.

A second example is a volcanic tunneling eruption [12]. Based on the ultimate surface deposits, it appears that a magma chamber under a certain volcano initially contained magma A. At a later time, a second, distinct magma B containing more water in solution entered the chamber, and a volcanic eruption ensued. Remarkably, almost pure magma B erupted out of the volcano before magma A was deposited on top of the magma B deposit. Magma B somehow tunneled through the magma chamber without entraining much of the resident magma A. Vesiculation bubbles from water coming out of solution in magma B reduced its average density as it rose through the magma chamber. The entrainment rate was reduced by the resulting buoyancy change, due to acceleration effects. Magma B was able to tunnel through magma A with little entrainment or mixing.

Compared to the ordinary buoyant plume, the increasing buoyancy from the latent heat release generates additional baroclinic torques. Their induced velocities conspire to narrow the width of the engulfment tongues into the plume. Since turbulent entrainment is dominated by these tongues [13], the entrainment in accelerating plumes is thereby reduced, in some cases dramatically [11] and [14]. For example, inside of thunderstorms, pure sea-level air can be transported up to the stratosphere essentially without dilution.

### III.

### MODEL

Turbidity currents are not subject to a phase change, but nonetheless have a potential source of additional buoyancy. Erosion of particles from the bed underneath a turbidity current tends to increase the (negative) buoyancy and hence  $g'$ . In turn, this tends to increase the Richardson number. In order to maintain  $Ri = 1$ , the

speed  $W$  tends to increase over what it would otherwise be, according to Eq. (1).

Increasing speed has two consequences. First, this permits additional erosion and continued suspension of bed particles into the flow. Second, the turbulent entrainment rate of pure, upper fluid into the current tends to be inhibited by acceleration effects.

Acceleration effects are quantified by a dimensionless acceleration parameter

$$\alpha = -d\tau_v/dt,$$

where  $\tau_v$  is the vortex rotation period [10]. For an unforced turbulent flow, the vortex rotation period is essentially equal to the chronological age of the vortex, and  $\alpha$  is just equal to its unforced value  $\alpha^*$

$$\alpha = \alpha^* < 0.$$

There may be acceleration effects even if the vortex rotation period is constant or increases only slowly in time. An example is the exponential jet, where the vortex rotation period is constant and equal to the e-folding time scale of the forcing [9]. The spreading angle is reduced by about 20% compared to the unforced jet.

The rate of bed erosion is a very strong function of the flow speed [15][16]. This is necessary for an erosion feedback mechanism to increase the buoyancy.

According to this model of turbidity currents, sediment can propagate much further over an erodible bed than a non-erodible one. The increased exchange of momentum with the erodible bed is predicted to be more than counterbalanced by the reduced exchange of momentum with the fluid above via entrainment.

In the extreme limit of precisely zero entrainment rate, a turbidity current flowing down a constant slope of uniform, erodible bed particles would never stop. Its thickness would remain constant forever, as it is accelerated hydraulically. In general, to the extent that acceleration effects reduce the entrainment rate, the current would travel farther than otherwise.

The model is based on the fact that acceleration only affects the entrainment rate when the change in vortex rotation rate is appreciable during one vortex rotation, the time scale of importance to the dynamics of the vortex. So a turbidity current would only experience

reduced entrainment from acceleration effects if the buoyancy continually increases. Once the normalized rate of increase in buoyancy falls below some threshold value, the acceleration effects are anticipated to disappear. The physics would revert back to the non-accelerating case, as observed in the turbulent jet [9].

Ambient fluid above the turbidity current is entrained at a rate  $v_a$ . In the limit of vanishing entrainment rate, the thickness of the turbidity current is approximately a constant,  $0$ . The vortex rotation period is

$$\tau_v = \delta_0/W = (\tau_0 - \alpha t) = \sqrt{(\delta_0/g')}.$$

Thus

$$W = \delta_0/(\tau_0 - \alpha t) \quad (3)$$

and

$$g' = \delta_0/(\tau_0 - \alpha t)^2. \quad (4)$$

If the Richardson number is unity, the time rate of change of  $g'$  due to erosion is

$$dg'/dt = 2\alpha\delta_0/(\tau_0 - \alpha t)^2 = 2\alpha W^3/\delta_0^2. \quad (5)$$

The bed erosion rate  $v_b$  is approximately proportional to the cube of the friction velocity [15][16], in accord with eq. (5).

According to this model, a turbidity current under the right conditions might approach an acceleration regime of vanishing entrainment. The velocity and the buoyant force would increase rapidly in time, according to Eq. (3) and (4). Even if the limit of vanishing entrainment is not reached, acceleration effects may partially inhibit entrainment, thereby increasing the speed and the propagation distance over that of a non-eroding density current.

A competing explanation is that the internal stratification within the turbidity current is responsible for inhibiting the entrainment. The particle concentration and/or size is greater near the bottom of the current, so that the flow is somewhat stratified [17]. Stratification is of course well known to inhibit turbulent entrainment [5].

Parker *et al.* [18] proposed a four-layer turbulence model that considers the energy of the flow under certain closure assumptions. Their model predicts the conditions under which the current will become “self-accelerating.” The present work suggests a physical mechanism for this transition, the dynamic effect of rotational acceleration on entrainment.

#### IV. CONCLUSIONS

A new model for the dynamics of turbidity currents is proposed. In analogy with phase change in clouds and in vesiculating magmas, the bed erosion under a turbidity current may increase the buoyant force sufficiently rapidly to induce acceleration effects such that the rate of entrainment rate of upper fluid into the current would be reduced. Ironically, bed erosion would increase the propagation distance of a turbidity current. Further work will attempt to model the physics numerically. Simple lab experiments with and without an erodible bed will also address the issue.

- 
1. E. Meiburg, J. McElwaine, and B. Kneller, “Turbidity currents and powder snow avalanches,” in *Handbook of Environmental Fluid Dynamics*, v. 1st ed. H. J. Fernando, Ch. 42, 557 (2013).
  2. [www.statista.com/statistics/377029/number-of-deaths-due-to-avalanches-in-the-us/](http://www.statista.com/statistics/377029/number-of-deaths-due-to-avalanches-in-the-us/)
  3. C. Cooper, J. Wood, and O. Andrieux, “Turbidity current measurements in the Congo Canyon,” OTC 23992, Offshore Technology Conference, Houston, 6-9 May, 2012 (2013).
  4. J. McElwaine, C. Cenedese, and J. Heninger, “Mixing in steady-state gravity currents,” 72nd Annual Meeting of the American Physical Society, Division of Fluid Dynamics, Seattle, (2019).
  5. J.S. Turner, *Buoyancy Effects in Fluids*, Cambridge (1973).
  6. B. Morton, G.I. Taylor, and J.S. Turner, “Turbulent gravitational convection from maintained and instantaneous sources,” *Proc. Roy. Soc. A*, **234** 1 (1956).
  7. R.E. Breidenthal, “The turbulent exponential jet,” *Phys. Fluids* **29** 2346 (1986).
  8. S.M. Kato, B.C. Groenewegen, and R.E. Breidenthal, “On turbulent mixing in nonsteady jets,” *AIAA J.* **25**(1) 165 (1987).
  9. Q. Zhang, and H. Johari, “Effects of acceleration on turbulent jets,” *Phys. Fluids* **8**(8) 2185 (1996).
  10. R.E. Breidenthal, “The effect of acceleration on turbulent entrainment,” *Phys. Scr.* **T132** 014001 (2008).
  11. G.S. Bhat and R. Narasimha, “A volumetrically heated jet: large-eddy structure and entrainment characteristics,” *J. Fluid Mech.* **325** 303 (1996).
  12. G.W. Bergantz and R.E. Breidenthal, “Non-stationary entrainment and tunneling eruptions: A dynamic link between eruption processes and magma mixing,” *Geoph. Res. Lett.* **28** 3075 (2001).
  13. A. Roshko, “Structure of turbulent shear flows: A new look,” *AIAA J.* **14**(10) 1349 (1976).
  14. R.E. Breidenthal, “Elements of entrainment,” in *Turbulencia, Escola, de Primavera em Transicao e Turbulencia*, ed. A.P. Silva Freire, A. Ilha, and R.E. Breidenthal, ABCM, Rio de Janeiro, **5**(2) 205 (2006).
  15. R.A. Bagnold, “The movement of desert sand,” *Proceedings of the Royal Society of London A* **157**(892), 594 (1936).
  16. M. Sekine, and K. Nishimori, “Erosion rate of cohesive sediment by running water,” ICSE - 4 *Fourth International Conference on Scour and Erosion*, Tokyo (2008).
  17. C.K. Paull, P.J. Talling, K.L. Maier, D. Parsons, J. Xu, D.W. Caress, R. Gwiazda, E.M. Lundsten, K. Anderson, J.P. Barry, M. Chaffey, T. O’Reilly, K.J. Rosenberger, J.A. Gales, B. Kieft, M. McGann, S.M. Simmons, S. McCann, E.J. Sumner, M.A. Claire, and M.J. Cartigny, “Powerful turbidity currents driven by dense basal layers,” *Nature Communications* **9** 4114 (2018).
  18. G. Parker, Y. Fukushima, and H.M. Pantin, “Self-accelerating turbidity currents,” *J. Fluid Mech.* **171**, 145 (1986).